

Routine Children's Immunization Schedule

Vaccine	2 months	4 months	6 months	12 months	18 months	4-6 years	Grade 6	Grade 9
DTaP-HB-IPV-Hib Diphtheria, tetanus, acellular pertussis, hepatitis B, polio, and Haemophilus influenzae B	✓	√	√					
DTaP-IPV-Hib Diphtheria, tetanus, acellular pertussis, polio, and Haemophilus influenzae B					✓			
DTaP-IPV/Tdap-IPV Diphtheria, tetanus, acellular pertussis, and polio						√		
Human Papillomavirus (HPV)							√ *	
Influenza (Flu)			√ **					
Measles, Mumps, and Rubella (MMR)				√		√		
Meningococcal C Conjugate (Men-C-C)	√	~		√				
Meningococcal Quadrivalent (Men-ACYW-135)								✓
Pneumococcal Conjugate	√	√	~	√				
Rotavirus	√	√	✓ ***					
Tdap Diphtheria, tetanus, and acellular pertussis								✓
Varicella				√		✓		

^{*} Two doses administered 6 months apart are required.

^{**} The flu shot should be administered every year. For those 6 months to 9 years old receiving the flu vaccine for the first time, 2 doses administered 4 weeks apart are required.

^{***} Rotavirus vaccine 3 dose schedule when using RotaTeq® vaccine

Children with specific medical conditions that place them at high risk of disease should receive an additional dose.